

# GETTING READY FOR THE CAPSULE PROCEDURE



## **Plan Ahead** Call **303-602-1926** if you have questions or need to schedule or change your appointment

- Your appointment is at Denver Health Outpatient Medical Center, Pavillion L, 660 N. Bannock St., 3rd floor. Patient/Visitor Parking is located behind the building.
- You will come to the GI Lab to take the capsule and then again to return the capsule once you have passed the capsule from your body and retrieved it.
- You will get 4 liters of Colyte for the preparation of the capsule procedure.

### **B** 3 days before the Capsule procedure

• Stop taking iron supplements.

#### The day before the Capsule procedure


- Stop taking antacids (Tums, Rolaids), Pepto-Bismol, and Carafate.
- You may eat a regular breakfast in the morning.
- 11 AM: DRINK ONLY CLEAR LIQUIDS for the rest of the day and evening.
  NO SOLID FOOD.
- Avoid any fluids that are red.
- **5-7 PM:** Drink 2 liters of the Colyte jug within the 2 hours this is a laxative and you will need to be near a bathroom.

### The day of the Capsule procedure

- 1 hour before your scheduled procedure, drink 1 liter of Colyte
- · After you finish the Colyte, do not eat or drink ANYTHING
- 2 hours after the capsule has been swallowed, drink the last 1 liter of Colyte. You may now resume drinking clear liquids.
- After 1 p.m., you can eat a light snack
- · Eat normally for dinner

# 8-72 hours after the Capsule procedure

- The capsule will typically take 8-48 hours to pass, if you haven't passed the capsule after 72 hours contact the GI Lab.
- Follow the instructions for getting your capsule once it is passed and return it to the GI Lab.



#### **GILAB**

Denver Health Outpatient Medical Center, Pavillion L 660 N. Bannock St. 3rd floor

303-602-1926